

Stress tensor sector of Holographic CFTs

Petar Tadić

Trinity College Dublin

R. Karlsson, M. Kulaxizi, A. Parnachev, PT
[1909.05775 \[hep-th\]](#)

R. Karlsson, M. Kulaxizi, A. Parnachev, PT
[2002.12254 \[hep-th\]](#)

HoloTube Junior

Introduction

AdS_{d+1}/CFT_d **correspondence:**

- ▶ Gravity in $(d + 1)$ -dimensional asymptotically Anti de-Sitter spacetime.
- ▶ Conformal Field Theory in d -dimensional spacetime.
- ▶ Weakly coupled gravity is dual to strongly coupled CFT.
- ▶ Can we use CFTs to define and describe quantum gravity?
- ▶ First, we need to understand the duality at classical level in gravity.
- ▶ Holographic CFTs ($C_T \rightarrow \infty$ and $\Delta_{\text{gap}} \rightarrow \infty$) have a weakly coupled gravity dual with local physics below the AdS scale.
[Heemskerk, Penedones, Polchinski, Sully, '09.]
- ▶ Second, learn more about strongly coupled CFTs.

Introduction

- ▶ We are interested in CFTs with large central charge $C_T \sim N^2 \rightarrow \infty$ and large gap $\Delta_{\text{gap}} \rightarrow \infty$. We study

$$\langle \mathcal{O}_H(\infty) \mathcal{O}_L(1) \mathcal{O}_L(z, \bar{z}) \mathcal{O}_H(0) \rangle,$$

where $\Delta_L \sim \mathcal{O}(1)$ and $\Delta_H \sim \mathcal{O}(C_T)$ while $\mu \sim \frac{\Delta_H}{C_T}$ is fixed and used as an expansion parameter.

- ▶ In CFT_2 there is an infinite-dimensional Virasoro algebra that strongly constraints correlators in $C_T \rightarrow \infty$ limit.
- ▶ $\langle \mathcal{O}_H(\infty) \mathcal{O}_L(1) \mathcal{O}_L(z) \mathcal{O}_H(0) \rangle \Big|_{\text{v.v.b.}} \sim e^{\Delta_L \mathcal{F}(\mu, z)}$,
where \mathcal{F} is a known function that can be expanded as $\mathcal{F}(\mu, z) = \sum_{k=0}^{\infty} \mu^k \mathcal{F}^{(k)}(z)$.

Introduction

- ▶ At $\mathcal{O}(\mu^k)$, $\mathcal{F}^{(k)}(z)$ can be written as

$$\mathcal{F}^{(k)}(z) = \sum_{\{i_p\}} b_{i_1 \dots i_k} f_{i_1}(z) \dots f_{i_k}(z), \quad \sum_{p=1}^k i_p = 2k,$$

where i_p are integers and

$$f_a(z) = (1-z)^a {}_2F_1(a, a, 2a, 1-z).$$

- ▶ $\mathcal{F}^{(k)}$ contains contributions from all quasi-primaries made of k stress-tensors:

$$\begin{aligned} \mathcal{F}^{(0)}(z) &\sim 1 && - && \hat{1}, \\ \mathcal{F}^{(1)}(z) &\sim b_2 f_2 && - && T(z), \\ \mathcal{F}^{(2)}(z) &\sim b_{22} f_2^2 + b_{13} f_1 f_3 && - && : T(z) \partial \dots \partial T(z) : \dots \end{aligned}$$

CFT_{d>2}

- ▶ **No** Virasoro symmetry anymore.
- ▶ We consider the exchange of multi-stress tensors.
- ▶ Their contribution to the correlator we denote as stress-tensor sector:

$$\langle \mathcal{O}_H(\infty) \mathcal{O}_L(1) \mathcal{O}_L(z, \bar{z}) \mathcal{O}_H(0) \rangle \Big|_{\text{multi-stress tensors}} = \sum_k \mu^k \mathcal{G}^{(k)}(z, \bar{z}).$$

- ▶ Now, multi-stress tensors are labeled by their spin s and twist τ ,

$$\tau = \Delta - s.$$

- ▶ Multi-stress tensors made with k stress-tensors contribute to $\mathcal{G}^{(k)}$.

Minimal-twist contributions in $\text{CFT}_{d>2}$

- ▶ First, we focus on minimal-twist multi-stress tensor operators.
- ▶ Operators with minimal twist, that contribute at $\mathcal{O}(\mu^k)$, can be schematically represented as:

$$[T^{(k)}]_{\tau,s} =: T_{\mu_1\nu_1} \cdots T_{\mu_{k-1}\nu_{k-1}} \partial_{\alpha_1} \cdots \partial_{\alpha_l} T_{\mu_k\nu_k} ;,$$

$$\tau = k(d-2),$$

$$s = 2k + l.$$

- ▶ The contribution of operators with the minimal twist is dominant over those from higher-twist operators in the lightcone limit $1 - \bar{z} \ll 1$.
- ▶ OPE coefficients of these operators are the same in all holographic CFTs. [Fitzpatrick, Huang, '19.][Fitzpatrick, Huang, Meltzer, Perlmutter, Simmons-Duffin, '20.]

Minimal-twist contributions in $\text{CFT}_{d>2}$

- ▶ In even-dimensional spacetime, we propose

$$\mathcal{G}^{(k)}(z, \bar{z}) \underset{\bar{z} \rightarrow 1}{\approx} \frac{(1 - \bar{z})^{k(\frac{d-2}{2})}}{[(1-z)(1-\bar{z})]^{\Delta_L}} \sum_{\{i_p\}} a_{i_1 \dots i_k} f_{i_1}(z) \dots f_{i_k}(z).$$

$$i_p \in \mathbb{N}, \quad \sum_{p=1}^k i_p = k \left(\frac{d+2}{2} \right) = \frac{\tau}{2} + s_{min},$$

where s_{min} is the minimal spin of operators that contribute at $\mathcal{O}(\mu^k)$ (For minimal-twist operators: $\tau = k(d-2)$, $s_{min} = 2k$).

- ▶ We now have to fix the unknown coefficients $a_{i_1 \dots i_k}$. These can be fixed via the lightcone bootstrap.

Lightcone bootstrap in CFT_4




Figure: Lightcone bootstrap - schematically

$$\mathcal{G}(z, \bar{z}) = \frac{1}{[(1-z)(1-\bar{z})]^{\Delta_L}} \sum_{\mathcal{O}_{\tau,s}} P_{\mathcal{O}_{\tau,s}}^{(HH,LL)} g_{\tau,s}^{(0,0)}(1-z, 1-\bar{z}),$$

where $\mathcal{O}_{\tau,s} \in \{1, T, : T \square^n \partial^l T :, \dots\}$.

$$\mathcal{G}(z, \bar{z}) = \frac{1}{(z\bar{z})^{\frac{1}{2}(\Delta_H + \Delta_L)}} \sum_{\mathcal{O}_{\tau',s'}} P_{\mathcal{O}_{\tau',s'}}^{(HL,HL)} g_{\tau',s'}^{(\Delta_{HL}, -\Delta_{HL})}(z, \bar{z}),$$

where $\mathcal{O}_{\tau',s'} \in \{:\mathcal{O}_H \square^n \partial^l \mathcal{O}_L :\}$.

Lightcone bootstrap in CFT_4

T-channel

- ▶ Conformal blocks of operators in the T-channel:

$$g_{\tau,s}^{(0,0)}(1-z, 1-\bar{z}) = (1-\bar{z})^{\frac{\tau}{2}} \left(f_{\frac{\tau}{2}+s}(z) + \mathcal{O}(1-\bar{z}) \right).$$

- ▶ At order $\mathcal{O}(\mu)$, there is only the stress-tensor contribution that is fixed by the Ward identity:

$$\begin{aligned} \mathcal{G}^{(1)}(z, \bar{z}) &= \frac{1}{((1-z)(1-\bar{z}))^{\Delta_L-1}} \frac{\Delta_L}{120(z-\bar{z})} \left(f_3(z) + f_3(\bar{z}) \right) \\ &\underset{\bar{z} \rightarrow 1}{\approx} \frac{(1-\bar{z})}{((1-z)(1-\bar{z}))^{\Delta_L}} \frac{\Delta_L}{120} f_3(z) \end{aligned}$$

Lightcone bootstrap in CFT_4

S-channel : $\mathcal{O}_H \partial^{2n} \partial^l \mathcal{O}_L$:

$$g_{\Delta_H + \Delta_L + 2n + \gamma, l}^{(\Delta_{HL}, -\Delta_{HL})}(z, \bar{z}) \underset{\Delta_H \rightarrow \infty}{\approx} \frac{(z\bar{z})^{\frac{1}{2}(\Delta_H + \Delta_L + 2n + \gamma)}}{\bar{z} - z} \left(\bar{z}^{l+1} - z^{l+1} \right)$$

$$\gamma_{n,l} = \sum_{k=1}^{\infty} \mu^k \gamma_{n,l}^{(k)} = \sum_{k=1}^{\infty} \mu^k \sum_{p=0}^{\infty} \frac{\gamma_n^{(k,p)}}{|k+p|},$$

$$P_{n,l}^{(HL,HL)} = P_{n,l}^{(HL,HL); \text{MFT}} \sum_{k=0}^{\infty} \mu^k P_{n,l}^{(HL,HL);(k)}, \quad P_{n,l}^{(HL,HL);(k)} = \sum_{p=0}^{\infty} \frac{P_n^{(k,p)}}{|k+p|}.$$

Lightcone bootstrap in CFT_4

- ▶ $f_i(z) = q_{i,1}(z) + \log(z)q_{i,2}(z)$, where $q_{i,1/2}(z)$ are rational functions.
- ▶ By matching $\mathcal{O}(\mu)$ from S-channel with T-channel (stress-tensor contribution), we fix $\gamma_n^{(1,\rho)}$ and $P_n^{(1,\rho)}$.
- ▶ At $\mathcal{O}(\mu^2)$, terms that contain $\log^2(z)$ in the S-channel are fixed by $\gamma_n^{(1,\rho)}$ and $P_n^{(1,\rho)}$.
- ▶ Generally, at $\mathcal{O}(\mu^k)$, terms that contain $\log^i(z)$, $2 \leq i \leq k$, in the S-channel are fixed by OPE data up to $\mathcal{O}(\mu^{k-1})$.

Lightcone bootstrap in CFT_4

- ▶ On the other hand, ansatzes in the T-channel are of the following form:

$$\mathcal{G}^{(k)}(z, \bar{z}) \underset{\bar{z} \rightarrow 1}{\approx} \frac{(1 - \bar{z})^k}{((1 - z)(1 - \bar{z}))^{\Delta_L}} \left(\tilde{q}_{k,z} \log^k(z) + \dots \right. \\ \left. + \tilde{q}_{1,z} \log(z) + \tilde{q}_{0,z} \right),$$

where $\tilde{q}_{i,z}$ are rational functions of z that depend on unknown coefficients $a_{i_1 \dots i_k}$.

- ▶ Therefore, we can match terms that behave as $\log^i(z)$ for $2 \leq i \leq k$, with those from the S-channel calculation, that are fixed in terms of OPE data at subleading order in μ .

Lightcone bootstrap in CFT_4

- ▶ At $\mathcal{O}(\mu^2)$, we match terms that contain $\log^2(z)$ from T- and S-channel.
- ▶ Matching these terms fixes coefficients the unknown coefficients a_{33} , a_{24} and a_{15} in the ansatz:

$$\mathcal{G}^{(2)}(z, \bar{z}) \underset{\bar{z} \rightarrow 1}{\approx} \frac{(1 - \bar{z})^2}{[(1 - z)(1 - \bar{z})]^{\Delta_L}} \left(\frac{\Delta_L}{28800(\Delta_L - 2)} \right) \times$$
$$\left\{ (\Delta_L - 4)(\Delta_L - 3)f_3^2(z) + \frac{15}{7}(\Delta_L - 8)f_2(z)f_4(z) \right.$$
$$\left. + \frac{40}{7}(\Delta_L + 1)f_1(z)f_5(z) \right\}$$

- ▶ By this means, we reproduce the correlator calculated in [\[Kulaxizi, Ng, Parnachev, '19.\]](#).
- ▶ Same method has been used calculate $\mathcal{O}(\mu^4)$ contributions in $d = 4$ and $\mathcal{O}(\mu^2)$ in $d = 6$. [\[Karlsson, Kulaxizi, Parnachev, PT, '19.\]](#)

Non-minimal twist - CFT_4

[Karlsson, Kulaxizi, Parnachev, PT, '20.]

- ▶ In $d = 4$, at $\mathcal{O}(\mu^2)$, minimal twist is $\tau = 4$, while first non-minimal twist double stress tensors have twist $\tau = 6$.
- ▶ These are two families of such operators:

$$: T_{\mu\alpha} \partial_{\lambda_1} \dots \partial_{\lambda_s} T^{\alpha}_{\nu} :, \quad \frac{\tau}{2} + s_{min} = 5,$$

$$: T_{\mu\nu} \partial_{\lambda_1} \dots \partial_{\lambda_s} \partial^2 T_{\rho\sigma} :, \quad \frac{\tau}{2} + s_{min} = 7.$$

- ▶ Now, we propose:

$$\mathcal{G}^{(2,1)}(z, \bar{z}) \underset{\bar{z} \rightarrow 1}{\propto} \frac{(1 - \bar{z})^3}{((1 - z)(1 - \bar{z}))^{\Delta_L}} \left(b_{14} f_1 f_4 + b_{23} f_2 f_3 \right. \\ \left. + c_{16} f_1 f_6 + c_{25} f_2 f_5 + c_{34} f_3 f_4 \right).$$

Non-minimal twist - CFT₄

[Karlsson, Kulaxizi, Parnachev, PT, '20.]

- ▶ Again, we use the lightcone bootstrap to fix the unknown coefficients.
- ▶ We look for terms proportional to the $\log^2(z)$ in the S-channel calculation.
- ▶ We have to keep subleading corrections to the S-channel OPE data in large-spin limit.
- ▶ We get $b_{23}, c_{16}, c_{25}, c_{34}$ in terms of Δ_L and b_{14} .
- ▶ b_{14} is the OPE coefficient of : $T_{\mu\alpha} T^{\alpha}_{\nu}$:.
- ▶ **Generally:** The lightcone bootstrap does not fix the OPE coefficients of operators with spin $s = 0, 2$.

Exponentiation and OPE coefficients

- ▶ It is shown that one can write the minimal-twist stress-tensor sector up to the $\mathcal{O}(\mu^4)$ in $d = 4$ as

$$\mathcal{G}(z, \bar{z}) \underset{\bar{z} \rightarrow 1}{\approx} \frac{1}{[(1-z)(1-\bar{z})]^{\Delta_L}} e^{\Delta_L \mathcal{F}(\mu; z, \bar{z})},$$

for some function \mathcal{F} which is a rational function of Δ_L and remains $\mathcal{O}(1)$ as $\Delta_L \rightarrow \infty$.

- ▶ One can use the following relation

$$f_a(z) f_b(z) = \sum_{m=0}^{\infty} p[a, b, m] f_{a+b+2m}(z)$$

to read off the OPE coefficients of minimal-twist multi-stress tensors from ansatzes with fixed coefficients.

Conclusion and future developments

- ▶ We found a method for the efficient calculation of the multi-stress tensor contributions.
- ▶ We confirm the universality of OPE coefficients of minimal-twist multi-stress tensors.
- ▶ We find that the minimal-twist contributions exponentiate in analogy with the Virasoro vacuum block.

In future:

- ▶ What happens with minimal-twist OPE coefficients in theory with finite Δ_{gap} ? [Fitzpatrick, Huang, Meltzer, Perlmutter, Simmons-Duffin, '20.], [20xx.xx - Karlsson, Kulaxizi, Parnachev, PT]
- ▶ Exploring the possibility of summing all minimal-twist contributions in a closed analytic form.

THANK YOU.